

Analytic Proof Systems for Gödel logic \mathbf{G}_3

M. Ultlog*

July 02, 2024

Abstract

We give sequent calculus, analytic tableaux, natural deduction, and clause translation systems for resolution for Gödel logic \mathbf{G}_3 .

1 Introduction

In this paper we present calculi for Gödel logic \mathbf{G}_3 [7]. \mathbf{G}_3 has three truth values $f, *, t$ (with t designated), connectives $\top, \perp, \neg, \supset, \wedge, \vee$, and quantifiers \forall, \exists . Its syntax and semantics is detailed in section 2.

We first present a 3-sided sequent calculus in section 3. The fundamental idea for many-sided sequent calculi for finite-valued logics goes back to Schröter [12], Rousseau [9], Takahashi [14]. We follow the method given by Baaz, Fermüller, and Zach [4] and Zach [15] for constructing inference rules. This guarantees that our system automatically has soundness and completeness theorems, cut-elimination theorem, mid-sequent theorem, and Maehara lemma (interpolation). For proofs of these results see [4, 15].

Signed tableau systems for finite-valued logics were proposed by Surma [13] and Carnielli [6], and generalized by Hähnle [8]. In section 4, we present a signed tableau system for Gödel logic.

Many-valued natural deduction systems for finite-valued logics have been investigated by Baaz, Fermüller, and Zach [3] and Zach [15]. We give the introduction and elimination rules for the natural deduction system for \mathbf{G}_3 in section 5.

In addition to Hähnle’s work on tableaux-based theorem proving for finite-valued logic, Baaz and Fermüller [1] have studied resolution calculi for clauses of signed literals. In order for these calculi to be used to prove that formulas of Gödel logic are tautologies or follow from some others, it is necessary to produce sets of signed clauses. In section 6, we present a translation calculus that yields a set of clauses from a set of formulas.

*Cite as: Multlog (2024), “Analytic proof systems for Gödel logic \mathbf{G}_3 .” [PDF generated by MULTLOG, v. 1.16a, <https://logic.at/multlog>]. Available at <https://logic.at/multlog/goedel.pdf>.

See appendices A and B for the specification of Gödel logic.

The rules we provide are optimal in each case, and use the algorithms developed by Salzer [10, 11].

2 Syntax and semantics

Definition 1. The *first order language* \mathcal{L} for Gödel logic consists of

1. free variables: a_0, a_1, a_2, \dots
2. bound variables: x_0, x_1, x_2, \dots
3. function symbols of arity i ($i \in \mathbb{N}$), including constants: $f_0^i, f_1^i, f_2^i, \dots$
4. predicate constants of arity i ($i \in \mathbb{N}$): $P_0^i, P_1^i, P_2^i, \dots$
5. propositional connectives, arity given in parenthesis: \top (0), \perp (0), \neg (1), \supset (2), \wedge (2) and \vee (2)
6. quantifiers: \forall and \exists
7. auxiliary symbols: “(”, “)” and “,”

Terms are defined inductively: Every individual constant and free variable is a term. If f^n is a function symbol of arity n , and t_1, \dots, t_n are terms, then $f^n(t_1, \dots, t_n)$ is a term.

Formulas are also defined inductively:

1. If P^n is a predicate symbol of arity n , and t_1, \dots, t_n are terms, then $P^n(t_1, \dots, t_n)$ is a formula. It is called *atomic* or an *atom*.
2. The propositional constant \top is an (atomic) formula.
3. The propositional constant \perp is an (atomic) formula.
4. If A is a formula, so is $\neg A$.
5. If A and B are formulas, so is $(A \supset B)$.
6. If A and B are formulas, so is $(A \wedge B)$.
7. If A and B are formulas, so is $(A \vee B)$.
8. If A is a formula not containing the bound variable x , a is a free variable and Q is a quantifier, then $(Qx)A(x)$, where $A(x)$ is obtained from A by replacing a by x at every occurrence of a in A , is a formula.

A formula is called *open*, if it contains free variables, and *closed* otherwise. A formula without quantifiers is called *quantifier-free*.

As a notational convention, lowercase letters from the beginning of the alphabet (a, b, c, \dots) will be used to denote free variables, f, g, h, \dots for function symbols and constants, x, y, z, \dots for bound variables, all possibly indexed. Uppercase letters A, B, C, \dots will stand for formulas, greek letters $\Gamma, \Delta, \Lambda, \dots$ for sequences and sets of formulas, t and s for terms. The symbols \Box and \mathbf{Q} stand for general propositional connectives and quantifiers, respectively.

Definition 2. We will use α as a variable for free variables (*eigenvariable*) and τ as a variable for terms (*term variable*). A formula consisting of some formula variables, eigenvariables and term variables is called a schema.

A *pre-instance* A' of a schema A is an actual formula from the formulas of \mathcal{L} which contains occurrences of the eigenvariables and term variables of A .

An *instance* A'' of A is a pre-instance A' of A , where the eigenvariables and term variables have been replaced by free variables and terms not occurring in A' .

Definition 3. The *matrix* for Gödel logic is given by:

1. the set of *truth values* $V = \{f, *, t\}$,
2. the set $V^+ = \{t\} \subseteq V$ of *designated truth values*,
3. the truth functions for connectives $\top, \perp, \neg, \supset, \wedge$ and \vee , as given below;
4. the truth functions for quantifiers \forall and \exists , as given below.

The set of *undesigned values* is $V^- = V \setminus V^+ = \{f, *\}$.

The truth functions for connectives $\top, \perp, \neg, \supset, \wedge$ and \vee are defined by

$\tilde{\top} \equiv t$	$\tilde{\perp} \equiv f$	<table style="border-collapse: collapse;"> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">$\tilde{\neg}$</td><td style="padding: 2px 5px;"></td></tr> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">f</td><td style="padding: 2px 5px;">t</td></tr> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">$*$</td><td style="padding: 2px 5px;">f</td></tr> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">t</td><td style="padding: 2px 5px;">f</td></tr> </table>	$\tilde{\neg}$		f	t	$*$	f	t	f	<table style="border-collapse: collapse;"> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">$\tilde{\supset}$</td><td style="padding: 2px 5px;">f</td><td style="padding: 2px 5px;">$*$</td><td style="padding: 2px 5px;">t</td></tr> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">f</td><td style="padding: 2px 5px;">t</td><td style="padding: 2px 5px;">t</td><td style="padding: 2px 5px;">t</td></tr> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">$*$</td><td style="padding: 2px 5px;">f</td><td style="padding: 2px 5px;">t</td><td style="padding: 2px 5px;">t</td></tr> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">t</td><td style="padding: 2px 5px;">f</td><td style="padding: 2px 5px;">$*$</td><td style="padding: 2px 5px;">t</td></tr> </table>	$\tilde{\supset}$	f	$*$	t	f	t	t	t	$*$	f	t	t	t	f	$*$	t	<table style="border-collapse: collapse;"> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">$\tilde{\wedge}$</td><td style="padding: 2px 5px;">f</td><td style="padding: 2px 5px;">$*$</td><td style="padding: 2px 5px;">t</td></tr> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">f</td><td style="padding: 2px 5px;">f</td><td style="padding: 2px 5px;">f</td><td style="padding: 2px 5px;">f</td></tr> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">$*$</td><td style="padding: 2px 5px;">f</td><td style="padding: 2px 5px;">$*$</td><td style="padding: 2px 5px;">$*$</td></tr> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">t</td><td style="padding: 2px 5px;">f</td><td style="padding: 2px 5px;">$*$</td><td style="padding: 2px 5px;">t</td></tr> </table>	$\tilde{\wedge}$	f	$*$	t	f	f	f	f	$*$	f	$*$	$*$	t	f	$*$	t
$\tilde{\neg}$																																												
f	t																																											
$*$	f																																											
t	f																																											
$\tilde{\supset}$	f	$*$	t																																									
f	t	t	t																																									
$*$	f	t	t																																									
t	f	$*$	t																																									
$\tilde{\wedge}$	f	$*$	t																																									
f	f	f	f																																									
$*$	f	$*$	$*$																																									
t	f	$*$	t																																									
		<table style="border-collapse: collapse;"> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">$\tilde{\forall}$</td><td style="padding: 2px 5px;">f</td><td style="padding: 2px 5px;">$*$</td><td style="padding: 2px 5px;">t</td></tr> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">f</td><td style="padding: 2px 5px;">f</td><td style="padding: 2px 5px;">$*$</td><td style="padding: 2px 5px;">t</td></tr> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">$*$</td><td style="padding: 2px 5px;">$*$</td><td style="padding: 2px 5px;">$*$</td><td style="padding: 2px 5px;">t</td></tr> <tr><td style="border-right: 1px solid black; padding: 2px 5px;">t</td><td style="padding: 2px 5px;">t</td><td style="padding: 2px 5px;">t</td><td style="padding: 2px 5px;">t</td></tr> </table>	$\tilde{\forall}$	f	$*$	t	f	f	$*$	t	$*$	$*$	$*$	t	t	t	t	t																										
$\tilde{\forall}$	f	$*$	t																																									
f	f	$*$	t																																									
$*$	$*$	$*$	t																																									
t	t	t	t																																									

The truth functions for quantifiers \forall and \exists are defined by

$\tilde{\forall}$		$\tilde{\exists}$	
$\{f, *, t\}$	f	$\{f, *, t\}$	t
$\{f, *\}$	f	$\{f, *\}$	$*$
$\{f, t\}$	f	$\{f, t\}$	t
$\{f\}$	f	$\{f\}$	f
$\{*, t\}$	$*$	$\{*, t\}$	t
$\{*\}$	$*$	$\{*\}$	$*$
$\{t\}$	t	$\{t\}$	t

Definition 4. A structure $\mathcal{M} = \langle D, \Phi_{\mathcal{M}} \rangle$ for a language \mathcal{L} (an \mathcal{L} -structure) consists of the following:

1. A nonempty set D , called the *domain* (elements of D are called *individuals*).
2. A mapping $\Phi_{\mathcal{M}}$ satisfying the following:
 - (a) Each free variable of \mathcal{L} is mapped to an element of D .
 - (b) Each n -ary function symbol f of \mathcal{L} is mapped to a function $f_{\mathcal{M}} : D^n \rightarrow D$, or to an element of D if $n = 0$. Additionally, $\Phi_{\mathcal{M}}$ maps elements of D to themselves.
 - (c) Each n -ary predicate symbol P of \mathcal{L} is mapped to a function $P_{\mathcal{M}} : D^n \rightarrow V$, or to a element of V if $n = 0$.

Definition 5. Let \mathcal{M} be an \mathcal{L} -structure. An *assignment* s is a mapping from the free variables of \mathcal{L} to individuals.

An *interpretation* $\mathfrak{I} = \langle \mathcal{M}, s \rangle$ is an \mathcal{L} -structure $\mathcal{M} = \langle D, \Phi_{\mathcal{M}} \rangle$ together with an assignment s . The mapping $\Phi_{\mathcal{M}}$ can be extended in the obvious way to a mapping $\Phi_{\mathfrak{I}}$ from terms to individuals:

1. If t is a free variable, then $\Phi_{\mathfrak{I}} := s(t)$.
2. If t is of the form $f(t_1, \dots, t_k)$, where f is a k -ary function symbol and t_1, \dots, t_k are terms, then $\Phi_{\mathfrak{I}} := f_{\mathcal{M}}(\Phi_{\mathfrak{I}}(t_1), \dots, \Phi_{\mathfrak{I}}(t_k))$.

Definition 6. Given an interpretation $\mathfrak{I} = \langle \mathcal{M}, s \rangle$, we define the *valuation* $\text{val}_{\mathfrak{I}}$ for formulas A to truth values as follows:

1. If A is atomic, $A = P(t_1, \dots, t_n)$, where P is an n -ary predicate symbol and t_1, \dots, t_n are terms, then let $\text{val}_{\mathfrak{I}}(A) = P_{\mathcal{M}}(\Phi_{\mathfrak{I}}(t_1), \dots, \Phi_{\mathfrak{I}}(t_n))$.
2. If $A = \square(A_1, \dots, A_n)$, where A_1, \dots, A_n are formulas, and $\tilde{\square}$ is the associated truth function to \square , then $\text{val}_{\mathfrak{I}}(A) = \tilde{\square}(\text{val}_{\mathfrak{I}}(A_1), \dots, \text{val}_{\mathfrak{I}}(A_n))$.
3. If $A = (\mathbb{Q}x)(B(x))$, and $\tilde{\mathbb{Q}}$ is the associated truth function to \mathbb{Q} , then $\text{val}_{\mathfrak{I}}(A) = \tilde{\mathbb{Q}}\{\text{val}_{\mathfrak{I}}(B(d)) \mid d \in D\}$.

Definition 7. An interpretation \mathfrak{I} *satisfies* a formula A , in symbols: $\mathfrak{I} \models A$, iff $\text{val}_{\mathfrak{I}}(A) \in V^+$.

Definition 8. Δ *entails* A iff $\mathfrak{I} \models A$ for every interpretation \mathfrak{I} such that $\mathfrak{I} \models B$ for all $B \in \Delta$. A is *valid* iff it is satisfied by every interpretation \mathfrak{I} .

3 Sequent calculus for Gödel logic

Definition 9 (Syntax of Sequents). A *sequent* Γ is a triple

$$\Gamma_f \mid \Gamma_* \mid \Gamma_t$$

of finite sequences Γ_v of formulas, where $v \in V$. The Γ_v are called the *components* of Γ .

For a sequence of formulas Δ , and $W \subseteq V$, let $[W:\Delta]$ denote the sequent whose component Γ_v is Δ if $v \in W$ and empty otherwise. For $\{w_1, \dots, w_k\}:\Delta$ we also write $[w_1, \dots, w_k:\Delta]$. If Γ and Γ' are sequents, then Γ, Γ' denotes the component-wise union, i.e., the v -component of Γ, Γ' is Γ_v, Γ'_v .

Definition 10. Let \mathfrak{J} be an interpretation. \mathfrak{J} *satisfies* a sequent Γ iff there is a $v \in V$ so that for some formula $A \in \Gamma_v$, $\text{val}_{\mathfrak{J}}(A) = v$. \mathfrak{J} is called a *model* of Γ , in symbols $\mathfrak{J} \models \Gamma$.

Γ is called *satisfiable* iff there is an interpretation \mathfrak{J} so that $\mathfrak{J} \models \Gamma$ and *valid* iff for every interpretation \mathfrak{J} , $\mathfrak{J} \models \Gamma$.

Proposition 11. $\Delta \models A$ iff the sequent $\Delta \mid \Delta \mid A$ is valid.

Definition 12. The *sequent calculus* for Gödel logic is given by:

1. axiom schemas of the form $A \mid A \mid A$,
2. weakening rules for every truth value v :

$$\frac{\Gamma}{\Gamma, [v:A]} \text{ W:v}$$

3. exchange rules for every truth value v :

$$\frac{\Gamma, [v:A, B], \Delta}{\Gamma, [v:B, A], \Delta} \text{ X:v}$$

4. contraction rules for every truth value v :

$$\frac{\Gamma, [v:A, A]}{\Gamma, [v:A]} \text{ C:v}$$

5. cut rules for every two truth values $v \neq w$:

$$\frac{\Gamma, [v:A] \quad \Delta, [w:A]}{\Gamma, \Delta} \text{ CUT:vw}$$

6. an introduction rule $\Box:v$ for every connective \Box and every truth value v , as specified below,
7. an introduction rule $\mathbf{Q}:v$ for every quantifier \mathbf{Q} and every truth value v , as specified below, where the free variables α occurring in the upper sequents satisfy the so-called *eigenvariable condition*: No α occurs in the lower sequent.

(2)–(5) are called *structural rules*. (6) and (7) are called *logical rules*.

The introduction rules for connective \top are given by

$$\frac{\Gamma_f \mid \Gamma_* \mid \Gamma_t}{\Gamma_f, \top \mid \Gamma_* \mid \Gamma_t} \top:f \quad \frac{\Gamma_f \mid \Gamma_* \mid \Gamma_t}{\Gamma_f \mid \Gamma_*, \top \mid \Gamma_t} \top:* \quad \frac{}{\Gamma_f \mid \Gamma_* \mid \Gamma_t, \top} \top:t$$

The introduction rules for connective \perp are given by

$$\frac{}{\Gamma_f, \perp \mid \Gamma_* \mid \Gamma_t} \perp:f \quad \frac{\Gamma_f \mid \Gamma_* \mid \Gamma_t}{\Gamma_f \mid \Gamma_*, \perp \mid \Gamma_t} \perp:* \quad \frac{\Gamma_f \mid \Gamma_* \mid \Gamma_t}{\Gamma_f \mid \Gamma_* \mid \Gamma_t, \perp} \perp:t$$

The introduction rules for connective \neg are given by

$$\frac{\Gamma_f \mid \Gamma_*, A \mid \Gamma_t, A}{\Gamma_f, \neg A \mid \Gamma_* \mid \Gamma_t} \neg:f \quad \frac{\Gamma_f \mid \Gamma_* \mid \Gamma_t}{\Gamma_f \mid \Gamma_*, \neg A \mid \Gamma_t} \neg:* \quad \frac{\Gamma_f, A \mid \Gamma_* \mid \Gamma_t}{\Gamma_f \mid \Gamma_* \mid \Gamma_t, \neg A} \neg:t$$

The introduction rules for connective \supset are given by

$$\frac{\Gamma_f, B \mid \Gamma_* \mid \Gamma_t \quad \Gamma_f \mid \Gamma_*, A \mid \Gamma_t, A}{\Gamma_f, A \supset B \mid \Gamma_* \mid \Gamma_t} \supset:f \quad \frac{\Gamma_f \mid \Gamma_*, B \mid \Gamma_t \quad \Gamma_f \mid \Gamma_* \mid \Gamma_t, A}{\Gamma_f \mid \Gamma_*, A \supset B \mid \Gamma_t} \supset:*$$

$$\frac{\Gamma_f, A \mid \Gamma_*, B \mid \Gamma_t, B \quad \Gamma_f, A \mid \Gamma_*, A \mid \Gamma_t, B}{\Gamma_f \mid \Gamma_* \mid \Gamma_t, A \supset B} \supset:t$$

The introduction rules for connective \wedge are given by

$$\frac{\Gamma_f, A, B \mid \Gamma_* \mid \Gamma_t}{\Gamma_f, A \wedge B \mid \Gamma_* \mid \Gamma_t} \wedge:f$$

$$\frac{\Gamma_f \mid \Gamma_*, B \mid \Gamma_t, B \quad \Gamma_f \mid \Gamma_*, A, B \mid \Gamma_t \quad \Gamma_f \mid \Gamma_*, A \mid \Gamma_t, A}{\Gamma_f \mid \Gamma_*, A \wedge B \mid \Gamma_t} \wedge:*$$

$$\frac{\Gamma_f \mid \Gamma_* \mid \Gamma_t, B \quad \Gamma_f \mid \Gamma_* \mid \Gamma_t, A}{\Gamma_f \mid \Gamma_* \mid \Gamma_t, A \wedge B} \wedge:t$$

The introduction rules for connective \vee are given by

$$\frac{\Gamma_f, B \mid \Gamma_* \mid \Gamma_t \quad \Gamma_f, A \mid \Gamma_* \mid \Gamma_t}{\Gamma_f, A \vee B \mid \Gamma_* \mid \Gamma_t} \vee:f$$

$$\frac{\Gamma_f, B \mid \Gamma_*, B \mid \Gamma_t \quad \Gamma_f \mid \Gamma_*, A, B \mid \Gamma_t \quad \Gamma_f, A \mid \Gamma_*, A \mid \Gamma_t}{\Gamma_f \mid \Gamma_*, A \vee B \mid \Gamma_t} \vee:*$$

$$\frac{\Gamma_f \mid \Gamma_* \mid \Gamma_t, A, B}{\Gamma_f \mid \Gamma_* \mid \Gamma_t, A \vee B} \vee:t$$

The introduction rules for quantifier \forall are given by

$$\frac{\Gamma_f, A(\tau) \mid \Gamma_* \mid \Gamma_t}{\Gamma_f, (\forall x)A(x) \mid \Gamma_* \mid \Gamma_t} \forall:f \quad \frac{\Gamma_f \mid \Gamma_*, A(\tau) \mid \Gamma_t \quad \Gamma_f \mid \Gamma_*, A(\alpha) \mid \Gamma_t, A(\alpha)}{\Gamma_f \mid \Gamma_*, (\forall x)A(x) \mid \Gamma_t} \forall:*$$

$$\frac{\Gamma_f \mid \Gamma_* \mid \Gamma_t, A(\alpha)}{\Gamma_f \mid \Gamma_* \mid \Gamma_t, (\forall x)A(x)} \forall:t$$

The introduction rules for quantifier \exists are given by

$$\frac{\Gamma_f, A(\alpha) \mid \Gamma_* \mid \Gamma_t}{\Gamma_f, (\exists x)A(x) \mid \Gamma_* \mid \Gamma_t} \exists:f \quad \frac{\Gamma_f \mid \Gamma_*, A(\tau) \mid \Gamma_t \quad \Gamma_f, A(\alpha) \mid \Gamma_*, A(\alpha) \mid \Gamma_t}{\Gamma_f \mid \Gamma_*, (\exists x)A(x) \mid \Gamma_t} \exists:*$$

$$\frac{\Gamma_f \mid \Gamma_* \mid \Gamma_t, A(\tau)}{\Gamma_f \mid \Gamma_* \mid \Gamma_t, (\exists x)A(x)} \exists:t$$

Definition 13. An upward tree of sequents is called a *proof* in the sequent calculus iff every leaf is an instance of an axiom, and all other sequents in it are obtained from the ones standing immediately above it by applying one of the rules. The sequent at the root of P is called its *end-sequent*. A sequent Γ is called *provable* iff it is the end-sequent of some proof.

Theorem 14 (Soundness and Completeness). *A sequent is provable if and only if it is valid.*

Proof. See Theorems 3.1 and 3.2 of Baaz et al. [4] or Theorems 3.3.8 and 3.3.10 of Zach [15]. \square

Corollary 15. *In Gödel logic, $\models A$ iff $\parallel A$ has a sequent proof, and $\Delta \models A$ iff $\Delta \mid \Delta \mid A$ has a proof.*

Theorem 16 (Cut-elimination). *The cut rule is eliminable in the sequent calculus for Gödel logic.*

Proof. See Theorem 4.1 of Baaz et al. [4] or Theorem 3.5.3 of Zach [15]. \square

Theorem 17 (Midsequent theorem). *The midsequent theorem holds in the sequent calculus for Gödel logic.*

Proof. See Theorem 6.1 of Baaz et al. [4] or Theorem 3.7.2 of Zach [15]. \square

Theorem 18 (Maehara lemma). *The Maehara lemma holds for the sequent calculus for Gödel logic.*

Proof. See Theorem 3.8.1 of Zach [15]. \square

4 Tableaux for Gödel logic

Although the method of Surma [13] and Carnielli [6] for obtaining signed analytic tableaux systems applies to Gödel logic, it has a drawback. As Hähnle [8] pointed out, to show that a formula is valid, it is required to provide as many closed tableaux as there are non-designated values. This is usually not desirable; the generalized approach by Hähnle [8] solves this problem. Below we give a tableau system for Gödel logic using the sets of signs $V \setminus \{v\}$, i.e., the tableau system exactly dual to that of Carnielli (in the sense of [2]).

Definition 19. A *signed formula* is an expression of the form $v:A$ where $v \in V$ and A is a formula.

Definition 20. A *tableau* for a set of signed formulas Δ is a downward rooted tree of signed formulas where each one is either an element of Δ or results from a signed formula in the branch above it by a branch expansion rule. A tableau is *closed* if every branch contains, for some formula A , the signed formulas $v: A$ for all $v \in V$, or a signed formula $v: A$ with a branch expansion rule that explicitly closes the branch (\otimes).

The branch expansion rules for connective \top are given by

$$\frac{t: \top}{\otimes}$$

The branch expansion rules for connective \perp are given by

$$\frac{f: \perp}{\otimes}$$

The branch expansion rules for connective \neg are given by

$$\frac{f: \neg A}{*: A} \quad \frac{t: \neg A}{f: A} \\ t: A$$

The branch expansion rules for connective \supset are given by

$$\frac{f: A \supset B}{f: B} \quad \frac{*: A}{*: A} \quad \frac{*: A \supset B}{*: B} \quad \frac{t: A}{t: A} \quad \frac{t: A \supset B}{f: A} \quad \frac{f: A}{f: A} \\ *: B \quad *: A \\ t: B \quad t: B$$

The branch expansion rules for connective \wedge are given by

$$\frac{f: A \wedge B}{f: A} \quad \frac{f: A \wedge B}{f: B} \quad \frac{*: A \wedge B}{*: B} \quad \frac{*: A \wedge B}{*: A} \quad \frac{t: A \wedge B}{t: B} \quad \frac{t: A \wedge B}{t: A}$$

The branch expansion rules for connective \vee are given by

$$\frac{f: A \vee B}{f: B} \quad \frac{f: A \vee B}{f: A} \quad \frac{*: A \vee B}{*: B} \quad \frac{*: A \vee B}{*: A} \quad \frac{f: A}{f: A} \quad \frac{t: A \vee B}{t: A} \\ *: B \quad *: A \\ t: B$$

The branch expansion rules for quantifier \forall are given by

$$\frac{f: (\forall x)A(x)}{f: A(\tau)} \quad \frac{*: (\forall x)A(x)}{*: A(\tau)} \quad \frac{*: (\forall x)A(x)}{*: A(\alpha)} \quad \frac{t: (\forall x)A(x)}{t: A(\alpha)} \\ t: A(\alpha)$$

The branch expansion rules for quantifier \exists are given by

$$\frac{f: (\exists x)A(x)}{f: A(\alpha)} \quad \frac{*: (\exists x)A(x)}{*: A(\tau)} \quad \frac{*: (\exists x)A(x)}{*: A(\alpha)} \quad \frac{t: (\exists x)A(x)}{t: A(\tau)} \\ f: A(\alpha)$$

Definition 21. An interpretation \mathfrak{I} *satisfies* a signed formula $v: A$ iff $\text{val}_{\mathfrak{I}}(A) \neq v$. A set of signed formulas is *satisfiable* if some interpretation \mathfrak{I} satisfies all signed formulas in it.

Theorem 22. *A set of signed formulas is unsatisfiable iff it has a closed tableau.*

Proof. Apply Theorems 4.14 and 4.21 of Hähnle [8]; interpreting $v: A$ as $S \ A$ where $S = V \setminus \{v\}$. \square

Corollary 23. *In Gödel logic, $\models A$ iff $\{v: A \mid v \in V^+\}$ has a closed tableau. $\Delta \models A$ iff $\{v: B \mid v \in V^-, B \in \Delta\} \cup \{v: A \mid v \in V^+\}$ has a closed tableau.*

5 Natural deduction for Gödel logic

Let Γ be a (set) sequent, $V^+ \subseteq V$ the set of *designated truth values*. The set of non-designated truth values is then $V^- = V \setminus V^+$. We divide the sequent Γ into its designated part Γ^+ and its non-designated part Γ^- in the obvious way:

$$\begin{aligned}\Gamma^+ &:= \langle \Gamma_v \mid v \in V^+ \rangle \\ \Gamma^- &:= \langle \Gamma_v \mid v \in V^- \rangle\end{aligned}$$

Definition 24. The *natural deduction rules* for Gödel logic are given by:

1. A weakening rule for all $v \in V^+$:

$$\frac{\Gamma^+}{\Gamma^+, [v: A]} \text{w}:v$$

2. For every connective \square and every truth value v an introduction rule $\square\text{I}:v$ (if $v \in V^+$) or an elimination rule $\square\text{E}:v$ (if $v \in V^-$).
3. For every quantifier Q and every truth value v an introduction rule $Q\text{I}:v$ (if $v \in V^+$) or an elimination rule $Q\text{E}:v$ (if $v \in V^-$).

The introduction and elimination rules for connective \top are given by

$$\frac{\frac{\Gamma_0^-, [[*: \top]] \quad \Gamma_1^-}{\Gamma_0^+, [t: \top] \quad \Gamma_1^+} \top\text{E}:f \quad \frac{\Gamma_0^-, [[f: \top]] \quad \Gamma_1^-}{\Gamma_0^+, [t: \top] \quad \Gamma_1^+} \top\text{E}:* \quad \overline{[t: \top]} \top\text{I}:t}$$

The introduction and elimination rules for connective \perp are given by

$$\frac{\Gamma_0^-, [[*: \perp]] \quad \Gamma_1^-}{\Gamma_0^+, [t: \perp] \quad \Gamma_1^+} \perp\text{E}:f \quad \frac{\Gamma_0^-, [[f: \perp]] \quad \Gamma_1^-}{\Gamma_0^+, [t: \perp] \quad \Gamma_1^+} \perp\text{E}:* \quad \frac{\Gamma_1^-}{\Gamma_1^+, [t: \perp]} \perp\text{I}:t$$

The introduction and elimination rules for connective \neg are given by

$$\begin{array}{c}
\frac{\Gamma_0^-, [[*: \neg A]] \quad \Gamma_1^-, [[*: A]]}{\Gamma_0^+, [t: \neg A] \quad \Gamma_1^+, [t: A]} \neg E: f \qquad \frac{\Gamma_0^-, [[f: \neg A]] \quad \Gamma_1^-}{\Gamma_0^+, [t: \neg A] \quad \Gamma_1^+} \neg E: * \\
\frac{\Gamma_1^-, [[f: A]]}{\Gamma_1^+, [t: \neg A]} \neg I: t
\end{array}$$

The introduction and elimination rules for connective \supset are given by

$$\begin{array}{c}
\frac{\Gamma_0^-, [[*: A \supset B]] \quad \Gamma_1^-, [[f: B]] \quad \Gamma_2^-, [[*: A]]}{\Gamma_0^+, [t: A \supset B] \quad \Gamma_1^+ \quad \Gamma_2^+, [t: A]} \supset E: f \\
\frac{\Gamma_0^-, [[f: A \supset B]] \quad \Gamma_1^-, [[*: B]] \quad \Gamma_2^-}{\Gamma_0^+, [t: A \supset B] \quad \Gamma_1^+ \quad \Gamma_2^+, [t: A]} \supset E: * \\
\frac{\Gamma_1^-, [[f: A]], [[*: B]] \quad \Gamma_2^-, [[f, *: A]]}{\Gamma_1^+, [t: B] \quad \Gamma_2^+, [t: B]} \supset I: t \\
\Gamma_1^+, \Gamma_2^+, [t: A \supset B]
\end{array}$$

The introduction and elimination rules for connective \wedge are given by

$$\begin{array}{c}
\frac{\Gamma_0^-, [[*: A \wedge B]] \quad \Gamma_1^-, [[f: A, B]]}{\Gamma_0^+, [t: A \wedge B] \quad \Gamma_1^+} \wedge E: f \\
\frac{\Gamma_0^-, [[f: A \wedge B]] \quad \Gamma_1^-, [[*: B]] \quad \Gamma_2^-, [[*: A, B]] \quad \Gamma_3^-, [[*: A]]}{\Gamma_0^+, [t: A \wedge B] \quad \Gamma_1^+, [t: B] \quad \Gamma_2^+ \quad \Gamma_3^+, [t: A]} \wedge E: * \\
\frac{\Gamma_1^- \quad \Gamma_2^-}{\Gamma_1^+, [t: B] \quad \Gamma_2^+, [t: A]} \wedge I: t \\
\Gamma_1^+, \Gamma_2^+, [t: A \wedge B]
\end{array}$$

The introduction and elimination rules for connective \vee are given by

$$\begin{array}{c}
\frac{\Gamma_0^-, [[*: A \vee B]] \quad \Gamma_1^-, [[f: B]] \quad \Gamma_2^-, [[f: A]]}{\Gamma_0^+, [t: A \vee B] \quad \Gamma_1^+ \quad \Gamma_2^+} \vee E: f \\
\frac{\Gamma_0^-, [[f: A \vee B]] \quad \Gamma_1^-, [[f, *: B]] \quad \Gamma_2^-, [[*: A, B]] \quad \Gamma_3^-, [[f, *: A]]}{\Gamma_0^+, [t: A \vee B] \quad \Gamma_1^+ \quad \Gamma_2^+ \quad \Gamma_3^+} \vee E: * \\
\Gamma_0^+, \dots, \Gamma_3^+
\end{array}$$

$$\frac{\Gamma_1^-}{\Gamma_1^+, [t: A, B]} \quad \forall I: t$$

The introduction and elimination rules for quantifier \forall are given by

$$\frac{\Gamma_0^-, [[*: (\forall x)A(x)]] \quad \Gamma_1^-, [[f: A(\tau)]]}{\Gamma_0^+, [t: (\forall x)A(x)] \quad \Gamma_1^+} \quad \forall E: f$$

$$\frac{\Gamma_0^-, [[f: (\forall x)A(x)]] \quad \Gamma_1^-, [[*: A(\tau)]] \quad \Gamma_2^-, [[*: A(\alpha)]]}{\Gamma_0^+, [t: (\forall x)A(x)] \quad \Gamma_1^+ \quad \Gamma_2^+, [t: A(\alpha)]} \quad \forall E: *$$

$$\frac{\Gamma_1^-}{\Gamma_1^+, [t: A(\alpha)]} \quad \forall I: t$$

The introduction and elimination rules for quantifier \exists are given by

$$\frac{\Gamma_0^-, [[*: (\exists x)A(x)]] \quad \Gamma_1^-, [[f: A(\alpha)]]}{\Gamma_0^+, [t: (\exists x)A(x)] \quad \Gamma_1^+} \quad \exists E: f$$

$$\frac{\Gamma_0^-, [[f: (\exists x)A(x)]] \quad \Gamma_1^-, [[*: A(\tau)]] \quad \Gamma_2^-, [[f, *: A(\alpha)]]}{\Gamma_0^+, [t: (\exists x)A(x)] \quad \Gamma_1^+ \quad \Gamma_2^+} \quad \exists E: *$$

$$\frac{\Gamma_1^-}{\Gamma_1^+, [t: A(\tau)]} \quad \exists I: t$$

Definition 25. A *natural deduction derivation* is defined inductively as follows:

1. Let A be any formula. Then

$$\frac{[V^-: A]}{[V^+: A]}$$

is a derivation of A from the assumption $[V^-: A]$ (an *initial derivation*).

2. If D_k are derivations of Γ_k^+, Δ_k^+ from the assumptions $\Gamma_k^-, \hat{\Delta}_k^-$, and

$$\frac{\left\langle \frac{\Gamma_k^-, [\Delta_k^-]}{\Gamma_k^+, \Delta_k^+} \right\rangle_{k \in K}}{\Pi^+}$$

is an instance of a deduction rule with $\hat{\Delta}_k^-$ a subsequence of Δ_k^- , and all eigenvariable conditions are satisfied, then

$$\frac{\langle D_k \rangle_{k \in K}}{\Pi^+}$$

is a derivation of Π^+ from the assumptions $\bigcup_{k \in K} \Gamma_k^-$. The formulas in $\hat{\Delta}_k^-$ which do not occur in $\bigcup_{k \in K} \Gamma_k^-$ are said to be *discharged* at this inference.

Theorem 26. *A partial sequent Γ^+ can be derived from the assumptions Γ^- in the natural deduction system for Gödel logic iff, for every interpretation \mathfrak{J} , either some formula in Γ_v^- ($v \in V^-$) evaluates to the truth value v , or there is a $w \in V^+$ and a formula in Γ_w^+ that evaluates to w .*

Proof. See Theorems 4.7 and 5.4 of Baaz et al. [3] or Theorems 4.2.8 and 4.3.4 of Zach [15]. \square

Corollary 27. $\Gamma \models A$ iff there is a natural deduction derivation of $[V^+:A]$ from $[V^-:\Gamma]$.

6 Resolution and clause formation rules for Gödel logic

The many-valued resolution calculus of Baaz and Fermüller [1] applies to Gödel logic. We present the framework here, as well as a language preserving clause translation system for Gödel logic.

Definition 28 (Signed formula). A *signed formula* is an expression of the form A^v , where A is a formula and $v \in V$. If A is atomic, A^v is a *signed atom*.

Definition 29 (Signed clause). A (signed) *clause* $C = \{A_1^{v_1}, \dots, A_n^{v_n}\}$ is a finite set of signed atoms (proper clause). The empty clause is denoted by \square .

An *extended clause* is a finite set of signed formulas.

Definition 30 (Semantics of clause sets). Let \mathcal{M} be a structure. \mathcal{M} *satisfies* a clause C iff for every assignment s , there is some signed formula $A^v \in C$, so that $\text{val}_{\mathfrak{J}}(A) = v$ (where $\mathfrak{J} = \langle \mathcal{M}, s \rangle$). \mathcal{M} *satisfies* a clause set \mathcal{C} iff it satisfies every clause in \mathcal{C} . \mathcal{C} is called *satisfiable* iff some structure satisfies it, and *unsatisfiable* otherwise.

The clause formation rules for connective \top are given by

$$\frac{\mathcal{C} \cup \{C \cup \{\top^f\}\}}{\mathcal{C} \cup \{C\}} \top:f \quad \frac{\mathcal{C} \cup \{C \cup \{\top^*\}\}}{\mathcal{C} \cup \{C\}} \top:* \quad \frac{\mathcal{C} \cup \{C \cup \{\top^t\}\}}{\mathcal{C}} \top:t$$

The clause formation rules for connective \perp are given by

$$\frac{\mathcal{C} \cup \{C \cup \{\perp^f\}\}}{\mathcal{C}} \perp:f \quad \frac{\mathcal{C} \cup \{C \cup \{\perp^*\}\}}{\mathcal{C} \cup \{C\}} \perp:* \quad \frac{\mathcal{C} \cup \{C \cup \{\perp^t\}\}}{\mathcal{C} \cup \{C\}} \perp:t$$

The clause formation rules for connective \neg are given by

$$\frac{\mathcal{C} \cup \{C \cup \{(\neg A)^f\}\}}{\mathcal{C} \cup \{C \cup \{A^*, A^t\}\}} \neg:f \quad \frac{\mathcal{C} \cup \{C \cup \{(\neg A)^*\}\}}{\mathcal{C} \cup \{C\}} \neg:*$$

$$\frac{\mathcal{C} \cup \{C \cup \{(\neg A)^t\}\}}{\mathcal{C} \cup \{C \cup \{A^f\}\}} \neg:t$$

The clause formation rules for connective \supset are given by

$$\frac{\mathcal{C} \cup \{C \cup \{(A \supset B)^f\}\}}{\mathcal{C} \cup \{C \cup \{B^f\}, C \cup \{A^*, A^t\}\}} \supset:f \quad \frac{\mathcal{C} \cup \{C \cup \{(A \supset B)^*\}\}}{\mathcal{C} \cup \{C \cup \{B^*\}, C \cup \{A^t\}\}} \supset:*$$

$$\frac{\mathcal{C} \cup \{C \cup \{(A \supset B)^t\}\}}{\mathcal{C} \cup \{C \cup \{A^f, B^*, B^t\}, C \cup \{A^f, A^*, B^t\}\}} \supset:t$$

The clause formation rules for connective \wedge are given by

$$\frac{\mathcal{C} \cup \{C \cup \{(A \wedge B)^f\}\}}{\mathcal{C} \cup \{C \cup \{A^f, B^f\}\}} \wedge:f$$

$$\frac{\mathcal{C} \cup \{C \cup \{(A \wedge B)^*\}\}}{\mathcal{C} \cup \{C \cup \{B^*, B^t\}, C \cup \{A^*, B^*\}, C \cup \{A^*, A^t\}\}} \wedge:*$$

$$\frac{\mathcal{C} \cup \{C \cup \{(A \wedge B)^t\}\}}{\mathcal{C} \cup \{C \cup \{B^t\}, C \cup \{A^t\}\}} \wedge:t$$

The clause formation rules for connective \vee are given by

$$\frac{\mathcal{C} \cup \{C \cup \{(A \vee B)^f\}\}}{\mathcal{C} \cup \{C \cup \{B^f\}, C \cup \{A^f\}\}} \vee:f$$

$$\frac{\mathcal{C} \cup \{C \cup \{(A \vee B)^*\}\}}{\mathcal{C} \cup \{C \cup \{B^f, B^*\}, C \cup \{A^*, B^*\}, C \cup \{A^f, A^*\}\}} \vee:*$$

$$\frac{\mathcal{C} \cup \{C \cup \{(A \vee B)^t\}\}}{\mathcal{C} \cup \{C \cup \{A^t, B^t\}\}} \vee:t$$

The clause formation rules for quantifier \forall are given by

$$\frac{\mathcal{C} \cup \{C \cup \{((\forall x)A(x))^f\}\}}{\mathcal{C} \cup \{C \cup \{A(f(\vec{a}))^f\}\}} \forall:f$$

$$\frac{\mathcal{C} \cup \{C \cup \{((\forall x)A(x))^*\}\}}{\mathcal{C} \cup \{C \cup \{A(f(\vec{a}))^*\}, C \cup \{A(b)^*, A(b)^t\}\}} \forall:*$$

$$\frac{\mathcal{C} \cup \{C \cup \{((\forall x)A(x))^t\}\}}{\mathcal{C} \cup \{C \cup \{A(b)^t\}\}} \forall:t$$

The clause formation rules for quantifier \exists are given by

$$\frac{\mathcal{C} \cup \{C \cup \{((\exists x)A(x))^f\}\}}{\mathcal{C} \cup \{C \cup \{A(b)^f\}\}} \exists:f$$

$$\frac{\mathcal{C} \cup \{C \cup \{((\exists x)A(x))^*\}\}}{\mathcal{C} \cup \{C \cup \{A(f(\vec{a}))^*\}, C \cup \{A(b)^*, A(b)^f\}\}} \exists:*$$

$$\frac{\mathcal{C} \cup \{C \cup \{((\exists x)A(x))^t\}\}}{\mathcal{C} \cup \{C \cup \{A(f(\vec{a}))^t\}\}} \exists:t$$

In the translation rules for quantifiers, the indicated free variables b are new free variables that do not already occur in the premise, and terms $f(\vec{a})$ are formed using a new function symbol f and \vec{a} all the free variables of the corresponding clause in the premise.

Theorem 31. *Let $T(\mathcal{C})$ be the result of exhaustively applying the translation rules to a clause set \mathcal{C} . Then $T(\mathcal{C})$ is a set of proper clauses, i.e., $T(\mathcal{C})$ contains only signed atoms (all connectives and quantifiers are eliminated). Furthermore, $T(\mathcal{C})$ is satisfiable iff \mathcal{C} is.*

Proposition 32. *Let A be a sentence and Δ be a set of sentences. Then*

1. $\models A$ iff $\{\{A^v \mid v \in V^-\}\}$ is unsatisfiable.
2. $\Delta \models A$ iff

$$\{\{B^w \mid w \in V^+\} \mid B \in \Delta\} \cup \{\{A^v \mid v \in V^-\}\}$$

is unsatisfiable.

Definition 33. A clause R is a *resolvent* of clauses C_1, C_2 if $R = (C_1 \setminus \{A_1^{v_1}\})\sigma \cup (C_2 \setminus \{A_2^{v_2}\})\sigma$ where

1. C_1 and C_2 have no free variables in common,
2. A_1 and A_2 are unifiable with most general unifier σ ,
3. $v_1 \neq v_2$.

If C_1 and C_2 have free variables in common, we say that R is a resolvent of C_1 and C_2 if it is a resolvent of variable-disjoint renamings C'_1 and C'_2 of C_1 and C_2 , respectively.

Definition 34. A *resolution refutation* of a clause set \mathcal{C} is a sequence of clauses C_1, \dots, C_n so that for every i , $C_i \in \mathcal{C}$ or C_i is a resolvent of C_j, C_k with $j, k < i$, and $C_n = \emptyset$.

Theorem 35. *A clause set \mathcal{C} is unsatisfiable iff it has a resolution refutation.*

Proof. See Theorems 3.14 and 3.19 of Baaz and Fermüller [1] or Theorems 2.5.5 and 2.5.8 of Zach [15]. \square

Corollary 36. $\Delta \models A$ iff

$$T(\{\{B^w \mid w \in V^+\} \mid B \in \Delta\} \cup \{\{A^v \mid v \in V^-\}\})$$

has a resolution refutation.

References

- [1] Matthias Baaz and Christian G. Fermüller. Resolution-based theorem proving for many-valued logics. *Journal of Symbolic Computation*, 19(4):353–391, 1995. DOI 10.1006/jSCO.1995.1021.
- [2] Matthias Baaz, Christian G. Fermüller, and Richard Zach. Dual systems of sequents and tableaux for many-valued logics. *Bulletin of the EATCS*, 51:192–197, 1993. DOI 10.11575/PRISM/38908.
- [3] Matthias Baaz, Christian G. Fermüller, and Richard Zach. Systematic construction of natural deduction systems for many-valued logics. In *23rd International Symposium on Multiple-Valued Logic. Proceedings*, pages 208–213. IEEE Press, 1993. DOI 10.1109/ISMVL.1993.289558.
- [4] Matthias Baaz, Christian G. Fermüller, and Richard Zach. Elimination of cuts in first-order finite-valued logics. *Journal of Information Processing and Cybernetics EIK*, 29(6):333–355, 1993. DOI 10.11575/PRISM/38801.
- [5] Matthias Baaz, Christian G. Fermüller, Gernot Salzer, and Richard Zach. Labeled calculi and finite-valued logics. *Studia Logica*, 61(1):7–33, 1998. DOI 10.1023/A:1005022012721.
- [6] Walter A. Carnielli. Systematization of finite many-valued logics through the method of tableaux. *The Journal of Symbolic Logic*, 52(2):473–493, 1987. DOI 10.2307/2274395.
- [7] Kurt Gödel. Zum intuitionistischen Aussagenkalkül. *Anzeiger der Akademie der Wissenschaften in Wien, Mathematisch-Naturwissenschaftliche Klasse*, 69:65–66, 1932.
- [8] Reiner Hähnle. *Automated Deduction in Multiple-Valued Logics*. Oxford University Press, 1993.
- [9] George Rousseau. Sequents in many valued logic I. *Fundamenta Mathematicae*, 60:23–33, 1967. URL <http://matwbn.icm.edu.pl/ksiazki/fm/fm60/fm6012.pdf>.
- [10] Gernot Salzer. MULTlog: An expert system for multiple-valued logics. In *Collegium Logicum*, pages 50–55. Springer, 1996. DOI 10.1007/978-3-7091-9461-4_3.

- [11] Gernot Salzer. Optimal axiomatizations of finitely valued logics. *Information and Computation*, 162(1-2):185–205, 2000. DOI 10.1006/inco.1999.2862.
- [12] Karl Schröter. Methoden zur Axiomatisierung beliebiger Aussagen- und Prädikatenkalküle. *Zeitschrift für mathematische Logik und Grundlagen der Mathematik*, 1(4):241–251, 1955. DOI 10.1002/malq.19550010402.
- [13] Stanislaw J. Surma. An algorithm for axiomatizing every finite logic. In David C. Rine, editor, *Computer Science and Multiple-Valued Logic: Theory and Applications*, pages 137–143. North-Holland, 1977.
- [14] Moto-o Takahashi. Many-valued logics of extended Gentzen style I. *Science Reports of the Tokyo Kyoiku Daigaku, Section A*, 9(231):271–292, 1968. URL <https://www.jstor.org/stable/43699119>.
- [15] Richard Zach. Proof theory of finite-valued logics. Diplomarbeit, Technische Universität Wien, 1993. DOI 10.11575/PRISM/38803.

A goedel.lgc – specification of Gödel logic

```

logic "G\"odel".
truth_values {f,*,t}.
designated_truth_values {t}.
ordering(linear, "f < * < t").
operator(true /0, mapping { t }).
operator(false/0, mapping { f }).
operator(neg /1, mapping { (t): f,
                          (*): f,
                          (f): t
                        }
        ).
operator(imp/2, table [
                      [ f, *, t,
                        f, t, t, t,
                        *, f, t, t,
                        t, f, *, t
                      ]
        ).
operator(and /2, inf(linear)).
operator(or /2, sup(linear)).
quantifier(forall, induced_by inf(linear)).
quantifier(exists, induced_by sup(linear)).

```

B goedel.cfg – L^AT_EX translations

```

% LaTeX configuration file for Goedel logic

texName(forall, \\forall).
texName(exists, \\exists).
texName(and, \\land ).

```



```
texName(or,    \\lor   ).
texName(neg,  \\lnot  ).
texName(imp,  \\supset).
texName(false, \\bot).
texName(true,  \\top).
texInfix(and).
texInfix(or).
texInfix(imp).
texPrefix(neg).

texExtra("ShortName", "\\mathbf{G}_3").
texExtra("Link", "https://logic.at/multlog/goedel.pdf").
texExtra("Preamble", "\\ESequentstrue").
```