Confluence without Termination via Parallel Critical Pairs

Bernhard Gramlich*

FB Informatik, Univ. Kaiserslautern, D-67653 Kaiserslautern, Germany
gramlich@informatik.uni-kl.de

Abstract. We present a new criterion for confluence of (possibly) non-terminating left-linear term rewriting systems. The criterion is based on certain strong joinability properties of *parallel critical pairs*. We show how this criterion relates to other well-known results, consider some special cases and discuss some possible extensions.

1 Introduction and Overview

For *abstract reduction (or abstract rewriting) systems* (ARSs for short) it is well-known that, under termination, confluence is equivalent to local confluence, via *Newman's Lemma*. For proving confluence of non-terminating ARSs, however, one usually needs much stronger local confluence properties. A very interesting unifying framework, based on so-called *decreasing diagrams*, for localizing confluence proofs (even without termination) in the general setting of *labelled abstract rewriting systems* has recently been developed by van Oostrom ([8]).

For *term rewriting systems* (TRSs for short), which are ARSs with some additional structure, local confluence can be characterized by confluence of *critical pairs* as expressed by the well-known *Critical Pair Lemma*. Hence, for (finite) terminating TRSs, this *critical pair test* yields decidability of confluence.

For non-terminating TRSs, however, the situation is much more difficult again. Even the absence of critical pairs does not guarantee confluence, as there exist non-terminating, non-overlapping TRSs which are (locally confluent but) not confluent (cf. e.g. [3]). These counterexamples must necessarily be non-left-linear.[2] In fact, TRSs which are left-linear and non-overlapping, i.e., *orthogonal*, are confluent (cf. e.g. [12]). This fundamentally important positive result has been considerably generalized by Huet ([3]) and further by Toyama ([13]) by allowing critical pairs, but imposing certain strong joinability properties on them (cf. Theorems 4, 6 and 8 below). Theorems 6 and 8 of Huet/Toyama are particularly interesting, since they do not require right-linearity. They are proved by showing strong confluence of *parallel reduction*, making essential use of the above mentioned particular joinability properties of (ordinary) critical pairs.

* This research was supported by 'Deutsche Forschungsgemeinschaft, SFB 314 D4'.

[2] Some interesting efforts for isolating the essence of the source of non-confluence in this case are described e.g. in [10], [15]. A decidable sufficient (syntactic) condition for confluence of non-left-linear, non-terminating TRSs is given in [15].

Our main new (and quite natural) idea now is that in order to ensure strong confluence of parallel reduction, one may also define and investigate the corresponding notion of *parallel critical pairs*. This concept indeed turns out to be very useful, since we are able to state and prove a new sufficient condition for strong confluence of parallel reduction (cf. Theorem 14) which is based on certain joinability properties for parallel critical pairs (cf. Definition 11).

Actually, the idea behind parallel critical pairs is not new. Implicitly, parallel critical pairs – or, more precisely, parallel critical peaks – are at the heart of so-called *critical pair criteria* for completion of terminating TRSs where certain (ordinary) critical pairs during completion can be ignored since they are redundant. Moreover, parallel critical pairs have also been used in another context for dealing with rewriting modulo some set of (non-orientable) equations ([11], [4]).

The rest of the paper is structured as follows. After introducing the necessary terminology, we present in Section 3 the known related results, give an example where none of the known confluence criteria applies, and motivate the introduction of parallel critical pairs. The main result of the paper, Theorem 14, is proved in Section 4. Its relation to the previous results, various illuminating examples and other related work are discussed in Section 5. Finally we conclude by discussing directions for further extending and generalizing our approach.

2 Preliminaries

We assume familiarity with the basic theory of *abstract reduction systems* as well as of the special case of *term rewriting systems*. For comprehensive surveys see e.g. [5], [2]. Furthermore we also use some basic facts about unification theory (cf. e.g. [1] for a recent survey).

2.1 Abstract Reduction Systems

An *abstract reduction system* (ARS) is a pair $\mathcal{A} = \langle A, \rightarrow \rangle$ consisting of a (base) set A and a binary relation $\rightarrow \subseteq A \times A$ also called *(abstract) reduction* or *(abstract) rewrite relation*. We use the standard notations $\rightarrow^+$, $\rightarrow^*$, $\rightarrow^=$ (or $\rightarrow^{\leq 1}$) for the transitive, transitive-reflexive and reflexive closure, respectively, of $\rightarrow$. The notations for the inverse relations are obtained by 'mirroring', e.g., we use $\leftarrow$ for the inverse of $\rightarrow$. Relation composition is denoted by $\circ$. Two elements $a, b \in A$ are said to be *joinable* (denoted by $a \downarrow b$) if there exists $c \in A : a \rightarrow^* c \ {}^*\!\leftarrow b$. If $a \rightarrow^* b$, we call b a *reduct* of a.

Definition 1. (confluence properties)
Let $\mathcal{A} = \langle A, \rightarrow \rangle$ be an ARS. Then $\mathcal{A}$ (or $\rightarrow$)

- is *confluent* (CONF) if ${}^*\!\leftarrow \circ \rightarrow^* \subseteq \rightarrow^* \circ {}^*\!\leftarrow$.
- is *Church-Rosser* (CR) if $\leftrightarrow^* \subseteq \rightarrow^* \circ {}^*\!\leftarrow$.
- is *locally confluent* or *weakly Church-Rosser* (WCR) if $\leftarrow \circ \rightarrow \subseteq \rightarrow^* \circ {}^*\!\leftarrow$.
- is *strongly confluent* (SCR) if $\leftarrow \circ \rightarrow \subseteq \rightarrow^* \circ {}^=\!\leftarrow$.
- is *subcommutative* (WCR$^{\leq 1}$) if $\leftarrow \circ \rightarrow \subseteq \rightarrow^= \circ {}^=\!\leftarrow$.

Confluence of an ARS $\mathcal{A} = \langle A, \to \rangle$ can be characterized in various equivalent ways. For instance, we have:

$$\mathrm{CR}(\to) \iff \mathrm{CONF}(\to) \iff \mathrm{CR}(\to^*) \iff \leftarrow \circ \to^* \subseteq \to^* \circ {}^*\leftarrow \ .$$

We note that due to the equivalence $\mathrm{CR}(\to) \iff \mathrm{CR}(\to^*)$ (by idempotence of $\cdot^*$), for proving confluence of $\to$ it suffices to prove confluence of $\rightsquigarrow$ for some reduction relation $\rightsquigarrow$ with the same transitive-reflexive closure, i.e. with $\rightsquigarrow^* = \to^*$. It is also well-known that confluence follows from strengthened versions of local confluence, according to the implications $\mathrm{WCR}^{\leq 1} \Longrightarrow \mathrm{SCR} \Longrightarrow \mathrm{CR}$. Local confluence (WCR), however, does in general not imply confluence (CR), but only under the additional assumption of termination.[3] This fundamentally important result, known as *Newman's Lemma*, is at the heart of many confluence proofs in the literature. Here, an ARS $\mathcal{A} = \langle A, \to \rangle$ (or $\to$) is said to be *terminating* if there exists no infinite reduction sequence $a_0 \to a_1 \to a_2 \to \ldots$ (of elements of A). $\mathcal{A}$ (or $\to$) is called *complete* (or *convergent*) if it is confluent and terminating.

2.2 Term Rewriting Systems

A term rewriting system $\mathcal{R}$ (over some set $\mathcal{T}(\mathcal{F}, \mathcal{V})$ of terms) may be viewed as the ARS $\langle \mathcal{T}(\mathcal{F}, \mathcal{V}), \to_{\mathcal{R}} \rangle$ where $\to_{\mathcal{R}}$ or simply $\to$ is the rewrite relation induced by the *rewrite rules* $l \to r \in \mathcal{R}$ ($l, r \in \mathcal{T}(\mathcal{F}, \mathcal{V})$ with $Var(r) \subseteq Var(l)$). A term is *linear* if no variable occurs more than once in it. A rewrite rule $l \to r$ is said to be *left-linear / right-linear* if l / r is linear. A TRS is *left-linear / right-linear* if all its rules are left-linear / right-linear.

For *substitutions* we use prefix notation, sometimes omitting parentheses, e.g., we write $\sigma(s)$ or σs to denote the result of applying the substitution σ to the term s. The *domain* and the *variable range* of a substitution σ are denoted by $Dom(\sigma)$ (or $Dom\ \sigma$) and $\mathcal{V}Ran(\sigma)$ (or $\mathcal{V}Ran\ \sigma$), respectively.

Positions (of terms) are also defined as usual, and denoted by p, q etc.. They are ordered by $\leq$ as usual. The 'empty' root position is denoted by λ. Concatenation of positions is denoted by juxtaposition. If $p \leq q$ we say that p is *above* q (or q is *below* p). For *subtracting* positions we use $\backslash$, i.e., we write $p \backslash q = q'$ if $p = qq'$. Two positions p and q are said to be *parallel* (or *independent, disjoint*), denoted by $p|q$, if neither $p \leq q$ nor $q \leq p$. These notations extend in a straightforward way to sets of positions.

The set of positions of a term s is denoted by $Pos(s)$. The sets of variable positions and of non-variable, i.e., function symbol, positions of s are denoted by $\mathcal{V}Pos(s)$ and $\mathcal{F}Pos(s)$, respectively. The subterm of s at some position $p \in Pos(s)$ is denoted by s/p. In order to indicate that t is the subterm of s at position $p \in Pos(s)$, we also write $s = s[p = t]$. The result of replacing in s the subterm at position $p \in Pos(s)$ by t is denoted by $s[p \leftarrow t]$. The result of (simultaneously) replacing in s the subterms at some set $P = \{p_1, \ldots, p_n\} \subseteq Pos(s)$ of parallel positions (of s) by $t_1, \ldots, t_n$, respectively, is denoted by $s[p_1 \leftarrow t_1] \ldots [p_n \leftarrow t_n]$ (where the order of the replacement is irrelevant) or simply $s[p_i \leftarrow t_i \mid 1 \leq i \leq n]$.

[3] A standard example for a locally confluent, but not (strongly) confluent (and hence necessarily non-terminating) system is $\{b \leftarrow a, b \to c, c \to b, c \to d\}$.

Similarly, $s[p_i = t_i \mid 1 \leq i \leq n]$ denotes s indicating additionally that $s/p_i = t_i$ for all i, $1 \leq i \leq n$.

If a term s rewrites to t at position $p \in Pos(s)$ using rule $l \to r$ and substitution σ, i.e., $s/p = \sigma l$, $t = s[p \leftarrow \sigma r]$, this is indicated by using the notation $s \to_{p,\sigma,l\to r} t$ or, when ignoring the applied rule and used substitution but keeping the position of the *contracted* (i.e., replaced) *redex* (i.e., reducible expression), by $s \to_p t$. Furthermore, we write $s \to_{\geq p} t$ if s reduces (in one step) to t by contracting some redex (in s) below $p \in Pos(s)$, and similarly for $s \to_{\leq p} t$.

A *unification problem* is a (finite) set (or multiset) of equations to be solved. Here we use $\doteq$ in order distinguish e.g. the singleton unification problem $s \doteq t$ from the (syntactic) equality $s = t$ (cf. [1] for further details).

For testing local confluence of TRSs the following notion originating from [6] is fundamental.

Definition 2. (critical pairs)
Let $\mathcal{R}$ be a TRS and let $l \to r, l' \to r'$ be two rules of $\mathcal{R}$ that w.l.o.g. do not have any common variable.[4] Suppose l/p, with $p \in \mathcal{F}Pos(l)$, and l' are unifiable with mgu σ, where $\mathcal{V}Ran(\sigma) \cap (Var(l) \setminus Var(l/p)) = \emptyset$.[5] Then the pair of reducts $\langle (\sigma l)[p \leftarrow \sigma r'], \sigma r \rangle$ is called a *critical pair* of $\mathcal{R}$, determined by *overlapping* $l' \to r'$ into $l \to r$ at position p. The divergence $(\sigma l)[p \leftarrow \sigma r'] \ {}_p\!\leftarrow \sigma l \to_\lambda \sigma r$ is the corresponding *critical peak*. If $p = \lambda$, we speak of an *outside critical peak* (or *critical overlay)*and *outside critical pair*, respectively. Otherwise, i.e., if $p > \lambda$, we speak of an *inside critical peak* and *inside critical pair*, respectively (following [13]). If the two rules are renamed versions of the same rule we do not consider the case $p = \lambda$ (which gives only rise to improper divergences). A critical pair $\langle s, t \rangle$ is said to be *joinable* if $s \downarrow t$. The set of all critical pairs between rules of $\mathcal{R}$ is denoted by $\mathrm{CP}(\mathcal{R})$. If $\mathrm{CP}(\mathcal{R}) = \emptyset$, $\mathcal{R}$ is said to be *non-overlapping*. It is called *orthogonal* if it is non-overlapping and left-linear.

Observe the asymmetry in the definition of critical pairs. This entails in particular, that for a critical overlay $t_1 \ {}_\lambda\!\leftarrow s \to_\lambda t_2$ we always get two corresponding critical pairs, namely $\langle t_1, t_2 \rangle$ and $\langle t_2, t_1 \rangle$. Moreover we note, that a critical pair may correspond to (i.e., be obtained from) several distinct critical peaks (if taking into account the position of the inside rewrite step and the applied rules).[6]

The *Critical Pair Lemma* ([3]) states that a TRS is locally confluent if and only if all its critical pairs are joinable. Combined with Newman's Lemma this yields ([6]): A terminating TRS is confluent (hence complete) if and only if all its critical pairs are joinable.

---

[4] Note that considering renamed versions of the same rule of $\mathcal{R}$ is allowed here, too.

[5] This variable condition ensures that the mgu σ does not introduce unnecessary identifications of variables in l. In case that the mgu of $t = l/p$ and l' is taken such that no new variables are introduced (which is always possible), this condition is vacuously satified.

[6] For the sake of readability we dispense here with a completely formal definition of critical peaks which is straightforward.

3 Known Results and Motivating Examples

Without termination, showing confluence of a TRS is much more difficult. For left-linear systems, a few results relying on strong confluence properties are known.

Definition 3. (strongly closed, [3]) A TRS $\mathcal{R}$ is called *strongly closed*, which we denote by SCCP($\mathcal{R}$),[7] if for every critical pair $\langle t_1, t_2 \rangle$ there exist terms t_3, t_4 such that $t_1 \to^* t_3 \,{}^{=}\!\leftarrow t_2$ and $t_1 \to^{=} t_4 \,{}^{*}\!\leftarrow t_2$.

Theorem 4. (strong confluence by strong closedness, [3]) *A left- and right-linear strongly closed TRS is strongly confluent, hence also confluent.*

The following very nice counterexample shows that right-linearity cannot be dropped in Theorem 4.

Example 1. **(due to J.-J. Lévy as cited in [3])** The TRS

$$\mathcal{R} = \left\{ \begin{array}{rcl} f(a,a) & \to & g(b,b) \\ a & \to & a' \\ f(a',x) & \to & f(x,x) \\ f(x,a') & \to & f(x,x) \\ g(b,b) & \to & f(a,a) \\ b & \to & b' \\ g(b'x) & \to & g(x,x) \\ g(x,b') & \to & g(x,x) \end{array} \right.$$

is left-linear, non-right-linear and non-terminating. Moreover, it is easily verified that $\mathcal{R}$ is strongly closed, but non-confluent. In fact, we have e.g. $f(a',a') \,{}^{*}\!\leftarrow f(a,a) \to^* g(b',b')$ but $f(a',a')$ and $g(b',b')$ do not have a common reduct.

For left-linear, but not necessarily right-linear TRSs one may use the following approach of Huet ([3]). Let $\mathcal{R}$ be a TRS. Then the *parallel reduction relation* (induced by $\mathcal{R}$), $\mathrel{-\!\!\!\!\!\!\rightarrow\!\!\!\!\!\!\rightarrow}_{\mathcal{R}}$ or simply $\mathrel{-\!\!\!\!\!\!\rightarrow\!\!\!\!\!\!\rightarrow}$, is defined as the smallest reflexive relation containing $\to$ and verifying

$$s_1 \mathrel{-\!\!\!\!\!\!\rightarrow\!\!\!\!\!\!\rightarrow} t_1 \wedge \ldots s_n \mathrel{-\!\!\!\!\!\!\rightarrow\!\!\!\!\!\!\rightarrow} t_n \implies f(s_1,\ldots,s_n) \mathrel{-\!\!\!\!\!\!\rightarrow\!\!\!\!\!\!\rightarrow} f(t_1,\ldots,t_n)$$

for all $f \in \mathcal{F}$ (n-ary). If s reduces to t by a parallel step contracting the redexes in s at some set $P = \{p_1,\ldots,p_k\}$ of parallel positions from P, this is also denoted by $s \mathrel{-\!\!\!\!\!\!\rightarrow\!\!\!\!\!\!\rightarrow}_P t$. Subsequently, we shall tacitly make use of some basic properties of parallel positions and parallel reduction, in particular the following ones:

- If $P_1, P_2 \subseteq Pos(s)$ are disjoint sets of positions of s such that all positions in $P_1 \uplus P_2$ are mutually disjoint, $t_p \to_\lambda t'_p$ for all $p \in P_1 \uplus P_2$, and $s = s[p = t_p \,|\, p \in P_1] \mathrel{-\!\!\!\!\!\!\rightarrow\!\!\!\!\!\!\rightarrow}_{P_1} s[p \leftarrow t'_p \,|\, p \in P_1] = s_1$, $s = s[p = t_p \,|\, p \in P_2] \mathrel{-\!\!\!\!\!\!\rightarrow\!\!\!\!\!\!\rightarrow}_{P_2} s[p \leftarrow t'_p \,|\, p \in P_2] = s_2$, then, defining $s_3 = s[p \leftarrow t'_p \,|\, p \in P_1 \uplus P_2]$, we get $s \mathrel{-\!\!\!\!\!\!\rightarrow\!\!\!\!\!\!\rightarrow}_{P_1} s_1 \mathrel{-\!\!\!\!\!\!\rightarrow\!\!\!\!\!\!\rightarrow}_{P_2} s_3$, $s \mathrel{-\!\!\!\!\!\!\rightarrow\!\!\!\!\!\!\rightarrow}_{P_2} s_2 \mathrel{-\!\!\!\!\!\!\rightarrow\!\!\!\!\!\!\rightarrow}_{P_1} s_3$, and $s \mathrel{-\!\!\!\!\!\!\rightarrow\!\!\!\!\!\!\rightarrow}_{P_1 \cup P_2} s_3$ (by commutation of the two independent parallel steps, and by combining both into one parallel step).

[7] This is to abbreviate 'strongly closed critical pairs'.

- If $p \leq q$ and $p'|p$, then $p'|q$.
- If P is a set of parallel positions of s, $q \in Pos(s)$ with $q \leq p$ for all $p \in P$, and s, t_1, t_2 coincide except for the respective subterms at position q, then $t_1 \ {}_P\!\!\leftarrow\!\!\!\!\!\!\parallel \ s \rightarrow_q t_2$ and $t_1/q \ {}_Q\!\!\leftarrow\!\!\!\!\!\!\parallel \ s/q \rightarrow_\lambda t_2/q$, where $Q = \{p \setminus q \,|\, p \in P\}$, are equivalent. In other words, the relevant part of a divergence $t_1 \ {}_P\!\!\leftarrow\!\!\!\!\!\!\parallel \ s \rightarrow_q t_2$ (with $q \leq p$ for all $p \in P$) can be obtained by extraction, stripping off the common context.

Clearly, parallel and ordinary reduction are related as follows: $\rightarrow \subseteq \!\!\!\!\!\!\parallel\!\!\rightarrow \subseteq \rightarrow^*$, hence also $\rightarrow^* = \!\!\!\!\!\!\parallel\!\!\rightarrow^*$. Thus, for showing confluence of $\rightarrow$, it suffices to prove (strong) confluence of $\!\!\!\!\!\!\parallel\!\!\rightarrow$.

Definition 5. (parallel closed, [3]) A TRS $\mathcal{R}$ is called *parallel closed* if for every critical pair $\langle t_1, t_2 \rangle$ of $\mathcal{R}$ we have: $t_1 \!\!\!\!\!\!\parallel\!\!\rightarrow t_2$.

Theorem 6. (subcommutativity of parallel reduction by parallel closedness, [3]) *If $\mathcal{R}$ is a left-linear parallel closed TRS, then parallel reduction (in $\mathcal{R}$) is subcommutative ($WCR^{\leq 1}(\!\!\!\!\!\!\parallel\!\!\rightarrow)$), hence $\mathcal{R}$ is confluent.*

Toyama ([13]) observed that, in Theorem 6 above, the condition that all critical pairs are parallel closed can be weakened without losing strong confluence of parallel reduction.

Definition 7. (critical pair condition of Huet/Toyama, [13])
We say that a TRS $\mathcal{R}$ satisfies the critical pair condition $\mathrm{CPC}_{\mathrm{HT}}$,[8] denoted by $\mathrm{CPC}_{\mathrm{HT}}(\mathcal{R})$, if for every critical peak $D: \ t_1 \ {}_p\!\!\leftarrow s \rightarrow_\lambda t_2$ of $\mathcal{R}$ (hence: $\langle t_1, t_2 \rangle \in \mathrm{CP}(\mathcal{R})$) we have:

(i) If $p = \lambda$, i.e., if D is an outside critical peak, then there exists a term t_3 such that $t_1 \!\!\!\!\!\!\parallel\!\!\rightarrow t_3 \ {}^*\!\!\leftarrow t_2$.[9]
(ii) If $p > \lambda$, i.e., if D is an inside critical peak, then $t_1 \!\!\!\!\!\!\parallel\!\!\rightarrow t_2$.

The conditions (i)-(ii) of Definition 7 are depicted in Figure 1 (dashed arrows in diagrams are existentially quantified).

Theorem 8. (strong confluence of parallel reduction, [13][10])
If $\mathcal{R}$ is a left-linear TRS satisfying the critical pair condition CPC_{HT}, then parallel reduction (in $\mathcal{R}$) is strongly confluent ($SCR(\!\!\!\!\!\!\parallel\!\!\rightarrow)$), hence $\mathcal{R}$ is confluent.

For motivating our approach to showing confluence via strong confluence of parallel reduction let us now consider the following modified version of Lévy's counterexample 1 above.

---

[8] Here, the subscript 'HT' is to abbreviate 'Huet/Toyama'.

[9] Note that due to the asymmetry in the definition of critical pairs / peaks this means that there must also exist a term t_4 with $t_1 \rightarrow^* t_4 \!\!\!\!\!\!\parallel\!\!\leftarrow t_2$!

[10] In fact, the corresponding result in [13], Corollary 3.2, p. 405, is obtained there by specializing a more general version, namely Theorem 3.1, p. 401, formulated in terms of a sufficient condition for *commutativity* of two left-linear TRSs.



Fig. 1. critical pair condition $\mathrm{CPC}_{\mathrm{HT}}$

Example 2. **(no known criterion applicable)**
The TRS

$$\mathcal{R} = \left\{ \begin{array}{r} f(a,a) \to b \\ a \to a' \\ f(a',x) \to f(x,x) \\ f(x,a') \to f(x,x) \\ f(a',a') \to b \\ b \to f(a',a') \end{array} \right.$$

is left-linear, but not right-linear, and non-terminating. Moreover, it is confluent which cannot be inferred from Theorem 8 since e.g. the inside critical peak $f(a',a) \leftarrow f(a,a) \to b$ is not parallel closed. Theorem 4 isn't applicable either, due to non-right-linearity of $\mathcal{R}$, though $\mathcal{R}$ is strongly closed. Nevertheless, parallel reduction (in $\mathcal{R}$) is strongly confluent (as will be shown). For instance, we observe that the divergence $f(a',a') \mathrel{|\!|\!\leftarrow} f(a,a) \to b$ can be 'strongly closed' via $f(a',a') \to b$ (or $b \to f(a',a')$). In Example 1 we had the divergence $f(a',a') \mathrel{|\!|\!\leftarrow} f(a,a) \to g(b,b)$ which could only be closed via $g(b,b) \to f(a,a) \mathrel{|\!|\!\rightarrow} f(a',a')$, but not 'strongly', i.e., there was no term t with $f(a',a') \to^* t \mathrel{|\!|\!\leftarrow} g(b,b)$. Furthermore we note that in both examples in the parallel step $f(a,a) \mathrel{|\!|\!\rightarrow} f(a',a')$ two parallel (inner) redexes are contracted both of which are critical w.r.t. the other outside step.

Intuitively, the observations above suggest to analyse more carefully local divergences where in one parallel step several (parallel) redex positions are critical w.r.t. the other step. From a more abstract point of view this means to study the following questions:

(1) What are *critical peaks / pairs* for the parallel reduction relation?
(2) Which conditions on the corresponding critical peaks / pairs for parallel reduction guarantee strong confluence?

For (1) we shall use the notion of *parallel critical peaks / pairs*. In fact, Definition 9 below only accounts for critical divergences of the form $t_1 \mathrel{|\!|\!\leftarrow} s \to_\lambda t_2$. However, this type of divergences (for parallel reduction) will turn out to be sufficient for characterizing strong confluence of $\mathrel{|\!|\!\rightarrow}$ (cf. Lemma 12 below). Concerning (2)

we shall present a *parallel critical pair condition* (cf. Definition 11) which indeed suffices for guaranteing strong confluence of $\twoheadrightarrow$ (cf. Theorem 14).

4 Main Result

Definition 9. (parallel critical pairs / peaks)
Let $\mathcal{R}$ be a TRS and let $l \to r, l_1 \to r_1, \ldots, l_n \to r_n$ be rules of $\mathcal{R}$ that w.l.o.g. do not have any common variable. Suppose $P = \{p_1, \ldots, p_n\}$ is a non-empty set of parallel non-variable positions of l. Suppose further that the unification problem $\{l_i \doteq l/p_i \mid 1 \leq i \leq n\}$ is solvable, let's say with mgu σ, where $\mathcal{V}Ran(\sigma) \cap (Var(l) \setminus (\bigcup_{i=1}^{n} Var(l/p_i))) = \emptyset$. Then the pair of reducts $\langle (\sigma l)[p_i \leftarrow \sigma r_i \mid p_i \in P], \sigma r \rangle$ is called a *parallel critical pair (of order n)* of $\mathcal{R}$, determined by *overlapping (in parallel)* the rules $l_i \to r_i$ into $l \to r$ at the (parallel) positions p_i (of l). The divergence $(\sigma l)[p_i \leftarrow \sigma r_i \mid 1 \leq i \leq n] \;{}_P\!\twoheadleftarrow \sigma l \to_\lambda \sigma r$ is the corresponding *parallel critical peak (of order n)*. If $n = 1$ and $l \to r$, $l_1 \to r_1$ are two renamed versions of the same rules, we do not consider the case $p = \lambda$ (which gives only rise to improper divergences). The set of all parallel critical pairs between rules of $\mathcal{R}$ is denoted by $\mathrm{PCP}(\mathcal{R})$. A parallel critical peak (and its corresponding parallel critical pair) is said to be *proper* if the inside parallel step in it contracts more than one redex (with the notation from above: $n \geq 2$).

In order to get some intuition for the introduced notion of parallel critical pairs / peaks and their computation let us give two simple examples.

Example 3. The TRS

$$\mathcal{R} = \begin{cases} f(g(x), h(x,y)) \to a \\ g(b) \to c \\ h(x,d) \to e \end{cases}$$

has two (parallel) critical peaks of order 1, $f(c, h(b,y)) \;{}_1\!\leftarrow f(g(b), h(b,y)) \to_\lambda a$ and $f(g(x), e) \;{}_2\!\leftarrow f(g(x), h(x,d)) \to_\lambda a$, obtained by overlapping the second and the third rule, respectively, into the first one. And there is one (proper) parallel critical peak of order 2, namely $f(c,e) \;{}_{\{1,2\}}\!\twoheadleftarrow f(g(b), h(b,d)) \to_\lambda a$, obtained either directly by overlapping in parallel the last two rules into the first, or recursively from both critical peaks of order 1 by an additional superposition. Hence, we have $\mathrm{PCP}(\mathcal{R}) = \{\langle f(c, h(b,y)), a\rangle, \langle f(g(x), e), a\rangle, \langle f(c,e), a\rangle\}$.

Example 4. The TRS

$$\mathcal{R} = \begin{cases} f(g(x), h(x)) \to a \\ g(b) \to d \\ h(c) \to d \end{cases}$$

has two (parallel) critical peaks of order 1, namely $f(d, h(b)) \;{}_1\!\leftarrow f(g(b), h(b)) \to_\lambda a$ and $f(g(c), d) \;{}_2\!\leftarrow f(g(c), h(c)) \to_\lambda a$, obtained by overlapping the second and the third rule, respectively, into the first one. However, there is no (proper) parallel critical peak of order 2, since simultaneously unifying $g(x)$ with $g(b)$ and $h(x)$ with $h(c)$ obviously fails. Hence, we get $\mathrm{PCP}(\mathcal{R}) = \{\langle f(d, h(b)), a\rangle, \langle f(g(c), d), a\rangle\}$.

Lemma 10. *Let $\mathcal{R}$ be a TRS. Then any divergence $t_1' \;{}_P\!\twoheadleftarrow s' \to_\lambda t_2'$, with $s' \to_{\lambda,\sigma,l\to r} t_2'$ and $s' \twoheadrightarrow_P t_1'$, P some set of n parallel non-variable positions p_i $(1 \leq i \leq n)$ from l, is an instance of a parallel critical peak $t_1 \;{}_P\!\twoheadleftarrow s \to_\lambda t_2$ of $\mathcal{R}$ (of order n).*

Proof. Routine. □

Next we introduce the central condition on parallel critical pairs that will enable us to prove strong confluence of parallel reduction.

Definition 11. (parallel critical pair condition)
We say that a TRS $\mathcal{R}$ satisfies the *parallel critical pair condition* PCPC (PCPC($\mathcal{R}$) for short) if for every parallel critical peak $D : t_1 \ {}_P\!\leftarrow\!\!\!

In fact, Lemma 12 can also be obtained as a special case from the following easy results about abstract reduction relations (taking $\to_1 = \mathrel{-\!\!\!\|\!\!\to}$ and $\to_2 = \to$).

Lemma 13. (relating strong confluence of related abstract reduction relations)
Let $\to_1$, $\to_2$ be abstract reduction relations (on some set A). Then the following properties hold:

(1) If ${}_1\!\leftarrow\circ\to_2 \subseteq \to_2^* \circ \overset{=}{}_1\!\leftarrow$ and $\to_1 \subseteq \to_2^* \subseteq \to_1^*$, then $SCR(\to_1)$.
(2) If $SCR(\to_1)$ and $\to_2 \subseteq \to_1^* \subseteq \to_2^*$, then ${}_1\!\leftarrow\circ\to_2 \subseteq \to_2^* \circ \overset{=}{}_1\!\leftarrow$.

Proof. Straightforward. □

Now we are prepared for the main result of the paper.

Theorem 14. (strong confluence of parallel reduction via parallel critical pairs) *If a left-linear TRS $\mathcal{R}$ satisfies the parallel critical pair condition PCPC, then parallel reduction (in $\mathcal{R}$) is strongly confluent (more succinctly: $PCPC(\mathcal{R}) \implies SCR(\mathrel{-\!\!\!\|\!\!\to})$), and hence, (ordinary reduction $\to$ in) $\mathcal{R}$ is also confluent.*

Proof. Let $\mathcal{R}$ be a left-linear TRS satisfying PCPC($\mathcal{R}$). According to Lemma 12 it suffices to show that whenever we have a local divergence of the form

$$D:\ t_1 \mathrel{\!\!\|\!\!\leftarrow} s \to t_2\ ,$$

there exists a term t_3 such that

$$t_1 \to^* t_3 \mathrel{\!\!\|\!\!\leftarrow} t_2\ .$$

Hence, suppose s reduces to t_1 by contracting n parallel redexes $s/p_1, \ldots, s/p_n$ in s using the rules $l_1 \to r_1, \ldots, l_n \to r_n$ of $\mathcal{R}$ and substitutions $\sigma_1, \ldots, \sigma_n$, respectively. We may assume that the set $P = \{p_1, \ldots, p_n\}$ of parallel redex positions is non-empty, i.e., $n \geq 1$ (the case $n = 0$ is trivial!). Further, suppose that s reduces to t_2 by contracting some redex s/q in s using some rule $l \to r$ of $\mathcal{R}$ with substitution σ. We proceed by a complete case distinction according to the relative positions of the contracted redexes (at $p_1, \ldots, p_n, q$ in s) in D (this case analysis is illustrated in Fig. 3 below).

(1) $q|P$ **(parallel redexes):** In this case we obviously get $t_1 \to_q t_3\ {}_P\!\mathrel{\!\!\|\!\!\leftarrow} t_2$ by commuting the steps, with t_3 defined by $t_3 = t_1[q \leftarrow t_2/q] = t_2[p_i \leftarrow t_1/p_i \mid 1 \leq i \leq n]$.
(2) $\exists p_i \in P : q = p_i$ **(overlay case):** If the applied rules at $s/p_i = s/q$, i.e., $l_i \to r_i$ and $l \to r$, are the same, we are done since we then have: $t_2 \mathrel{-\!\!\!\|\!\!\to}_{P\setminus\{p_i\}} t_1 =: t_3$. Otherwise, the local divergence $t_1/q\ {}_{\lambda,l_i\to r_i}\!\leftarrow s/q \to_{q,l\to r} t_2/q$ (obtained from D by considering only the subterm s/q) is an instance of a critical overlay (between the rules $l_i \to r_i$ and $l \to r$), hence by (the symmetric version of) the assumption PCPC($\mathcal{R}$)(i) we get $t_1/q \to^* t_3' \mathrel{\!\!\|\!\!\leftarrow} t_2/q$ for some t_3'. Combining the reductions appropriately and exploiting commutation of parallel redex contractions we obtain the existence of t_1', t_3 with $s \to_q s' \to^*_{\geq q} t_1', t_2 \mathrel{-\!\!\!\|\!\!\to}_{\geq q} t_1' \mathrel{-\!\!\!\|\!\!\to}_{P\setminus\{q\}} t_3$, $s \to_q s' \mathrel{-\!\!\!\|\!\!\to}_{P\setminus\{q\}} t_1 \to^*_{\geq q} t_3$. Hence, $t_1 \to^* t_3 \mathrel{\!\!\|\!\!\leftarrow} t_2$ as desired.



Fig. 3. illustration of the cases of the proof of Theorem 14

(3) $\exists p_i \in P : q > p_i$ **(the single redex is below one of the parallel redexes):** In this case we further distinguish between a variable and critical overlap.

(3.1) $\exists p \in \mathcal{V}Pos(l_i) : q \geq p_i p$ **(variable overlap below p_i):** Let $l_i/p = x \in \mathcal{V}$ and $p' = q \setminus (p_i p)$. Define σ_i' by $Dom\ \sigma_i' = Dom\ \sigma$, $\sigma_i' y = \sigma_i y$ for $y \neq x$, and $\sigma_i' x = \sigma_i x[p' \leftarrow \sigma r]$. Then we get $s = s[p_i = \sigma_i l_i] \mathbin{-\!\!\!\!\!\!\Vdash}_P t_1 = t_1[p_i = \sigma_i r_i] \mathbin{-\!\!\!\!\!\!\Vdash} t_1[p_i \leftarrow \sigma_i' r_i] =: t_3$ and $s \rightarrow_q t_2 \mathbin{-\!\!\!\!\!\!\Vdash}_P t_3$ as desired.

(3.2) $q \setminus p \in \mathcal{F}Pos(l_i)$ **(critical overlap below p_i):** Let $p' = q \setminus p_i$. Then the mirrored version of the (extracted) divergence $t_1/p_i = \sigma_i r_i \ {}_\lambda\!\leftarrow \sigma_i l_i =$

$s/p_i = s/p_i[p' = \sigma l] \to_{p'} s/p_i[p' \leftarrow \sigma r] = t_2/p_i$ is an instance of an (ordinary) inside critical peak of $\mathcal{R}$ (obtained by overlapping $l \to r$ into $l_i \to r_i$ at position p' in l_i). By assumption (ii) of PCPC($\mathcal{R}$) we obtain $t_1/p_i \to^*_{\geq\lambda} t_3'$ and $t_2/p_i \twoheadrightarrow_{\geq\lambda} t_3'$ for some t_3'. Combining the reductions appropriately and exploiting commutation of parallel redex contractions we obtain the existence of s', t_2', t_3 with $s \to_{p_i} s' \twoheadrightarrow_{P\setminus\{p_i\}} t_1 \to^*_{\geq p_i} t_3$, $s \to_q t_2 \twoheadrightarrow_{\geq p_i} t_2' \twoheadrightarrow_{P\setminus\{p_i\}} t_3$, hence $t_1 \to^* t_3 \mathrel{+\!\!\!

i.e., we do not have $f(a',a) \mathrel{-\!\!\!\|\!\!\!\rightarrow} b$. Moreover, the strongly closed criterion SCCP is also not applicable, because $\mathcal{R}$ is not right-linear.

Example 6. **(left- and right-linear case: PCPC $\not\Rightarrow$ CPC$_{\mathrm{HT}}$, SCCP)**
The TRS

$$\mathcal{R} = \begin{cases} f(a) \to f(g(b,b)) \\ a \to g(c,c) \\ c \to d \\ d \to b \\ b \to d \end{cases}$$

is left- and right-linear (and non-terminating). Moreover, it is easily checked that $\mathcal{R}$ satisfies PCPC($\mathcal{R}$), but neither CPC$_{\mathrm{HT}}$($\mathcal{R}$) nor SCCP($\mathcal{R}$).

Example 7. **(left- and right-linear case: SCCP $\not\Rightarrow$ PCPC)**
The TRS

$$\mathcal{R} = \begin{cases} f(a,b) \to c \\ a \to a' \\ b \to b' \\ c \to f(a',b) \\ c \to f(a,b') \\ c \to f(a,b) \end{cases}$$

is left- and right-linear (and non-terminating). Furthermore, as it is easy to check, we have SCCP($\mathcal{R}$) but neither CPC$_{\mathrm{HT}}$($\mathcal{R}$) nor PCPC($\mathcal{R}$). This means that $\mathcal{R}$ is strongly confluent. Note, however, that parallel reduction is only confluent here, but not strongly confluent.

Example 8. **(left-linear case: CPC$_{\mathrm{HT}}$ $\not\Rightarrow$ PCPC)**
The TRS

$$\mathcal{R} = \begin{cases} f(a,a) \to g(f(a,a)) \\ a \to b \\ f(b,x) \to g(f(x,x)) \\ f(x,b) \to g(f(x,x)) \end{cases}$$

is left-linear (and non-terminating). Furthermore, it is obviously parallel closed, i.e., CPC$_{\mathrm{HT}}$($\mathcal{R}$) holds. However, it doesn't satisfy the parallel critical pair condition PCPC($\mathcal{R}$) (because the proper parallel critical pair $\langle f(b,b), g(f(a,a))\rangle$ is not joinable from left to right, i.e., $f(b,b) \to^* g(f(a,a))$ does not hold).

The last example shows in particular, that CPC$_{\mathrm{HT}}$ is not subsumed in general by PCPC. However, there is one special case where CPC$_{\mathrm{HT}}$ is indeed (properly) subsumed by PCPC, namely when there are no proper parallel critical pairs (but only ordinary inside ones). In that case, parallel closedness of all inside critical pairs implies conditions (ii) and (iii) of PCPC, according to CPC$_{\mathrm{HT}}$(ii). This is obvious for PCPC(ii), and PCPC(iii) only has to hold for (ordinary) inside critical peaks which is guaranteed again by CPC$_{\mathrm{HT}}$(ii).

Finally, let us briefly discuss related work and possible extensions of our approach. First of all, it seems rather straightforward to generalize our approach to a 'commutation setting' analogous to the one of Toyama ([13]) mentioned above for Theorem 8.

Furthermore, we would like to mention one possible source of refinements for Theorem 14 (and related critical pair based confluence criteria) which is based

on making use of the ambiguity of rewriting. More precisely, it may be the case that a parallel critical peak $t_1 \ {}_{>\lambda}\!\leftarrow\!\!\!\!\!\!\parallel\ s \rightarrow_\lambda t_2$ as in PCPC(iii) is not joinable from left to right as required. However, it might e.g. be (an instance of) another outside critical peak $t_1' \ {}_{\lambda}\!\leftarrow s' \rightarrow_\lambda t_2'$ which satisfies PCPC(i), i.e., $t_1' \!\!\!\!\!\!\parallel\!\!\rightarrow t_3 \ {}^{*}\!\leftarrow t_2'$ and $t_1' \rightarrow^* t_4 \!\leftarrow\!\!\!\!\!\!\parallel\ t_2'$ for some t_3', t_4'. In this case the reasoning in the proof of Theorem 14 still applies. The following simple example ilustrates this kind of refinement.

Example 9. **(refining PCPC)**
The TRS

$$\mathcal{R} = \begin{cases} f(a) \rightarrow f(f(a)) \\ a \rightarrow b \\ f(x) \rightarrow f(b) \end{cases}$$

does not satisfy PCPC(iii) since for the inside critical peak $f(b) \ {}_{>\lambda}\!\leftarrow f(a) \rightarrow_\lambda f(f(a))$ we do not have $f(b) \rightarrow^* f(f(a))$. However, we also have the outside critical peak $f(b) \ {}_{\lambda}\!\leftarrow f(a) \rightarrow_\lambda f(f(a))$, which is strongly joinable in the sense of PCPC(i), due to $f(f(a)) \rightarrow f(b)$. Hence, parallel reduction in $\mathcal{R}$ is strongly confluent.

Similar refinements are also possible for Theorem 8.

Another direction of extending our approach is to combine it with the recent results of van Oostrom ([9]) who showed that the approach of Huet/Toyama can be generalized by replacing parallel reduction steps by the more general so-called *(complete) developments*, using essentially the same proof structure as in Theorem 8.[12] The basic idea of *developments* roughly is that a set of rewrite steps is considered to be *parallel* if a 'parallel extraction' of all the steps in the set is possible ([9]). This implies for instance, that in a (left-linear) TRS a derivation $s = \sigma l \rightarrow_p \sigma' l \rightarrow_{\lambda,\sigma',l\rightarrow r} \sigma' r$ (with $p \geq q$ for some $q \in \mathcal{V}Pos(l)$) can be viewed as a single parallel step. It seems that our approach via parallel critical pairs and the approach of van Oostrom generalizing the notion of being 'parallel' are 'orthogonal' to each other, hence it should be possible to combine both as well as the corresponding proof structures.

Similarly, it is conceivable that our approach can also be combined with (proof techniques for) confluence results for *higher-order rewrite systems* ([7]).

Last but not least we think that the notion of parallel critical pairs might turn out to be fruitful for other purposes, too. But this remains to be seen.

Acknowledgements: I thank Vincent van Oostrom and Claus-Peter Wirth for useful comments, Paul Taylor for his diagrams, and the anonymous referees for some constructive hints (in particular, the pointers to [11], [4]).

Final Note: It should be mentioned that Yoshihito Toyama, after having read (a longer version of) this paper, informed the author ([14]) about the existence of another very early paper, namely *"Y. Toyama: On the Church-Rosser property of term rewriting systems, NTT ECL Technical Report 17672 (Dec. 23, 1981), in Japanese"*, containing very similar ideas and results. The relationships between

[12] Actually, the corresponding result in [9] is even more general, since it also holds for (certain) left-linear higher-order systems, called *patterm rewriting systems* there.

the latter paper and ours as well as possible extensions and refinements will be investigated in joint future research.

References

1. F. Baader and J. Siekmann. Unification theory. In D. Gabbay, C. Hogger, and J. Robinson, eds., *Handbook of Logic in Artificial Intelligence and Logic Programming*, vol. 2 of *Handbooks of Logic in Computer Science and of Logic in Artificial Intelligence and Logic Programming*, pp. 41–126. Clarendon Press, Oxford, 1994.
2. N. Dershowitz and J.-P. Jouannaud. Rewrite systems. In J. van Leeuwen, ed., *Formal models and semantics, Handbook of Theoretical Computer Science*, vol. B, ch. 6, pp. 243–320. Elsevier - The MIT Press, 1990.
3. G. Huet. Confluent reductions: Abstract properties and applications to term rewriting systems. *Journal of the ACM*, 27(4):797–821, Oct 1980.
4. J.-P. Jouannaud, H. Kirchner, and J.-L. Rémy. Church-Rosser properties of weakly terminating term rewriting systems. In *Proc. 8th IJCAI*, vol. 2, pp. 909–915, Aug. 1983.
5. J. W. Klop. Term rewriting systems. In S. Abramsky, D. Gabbay, and T. Maibaum, eds., *Handbook of Logic in Computer Science*, vol. 2, ch. 1, pp. 2–117. Clarendon Press, Oxford, 1992.
6. D. E. Knuth and P. Bendix. Simple word problems in universal algebra. In J. Leech, editor, *Computational Problems in Abstract Algebra*, pages 263–297. Pergamon Press, Oxford, U. K., 1970. Reprinted 1983 in "Automation of Reasoning 2", Springer-Verlag, Berlin, pp. 342–376.
7. R. Mayr and T. Nipkow. Higher-order rewrite systems and their confluence. Technical Report TUM-I9433, TU München, Aug. 1994.
8. V. Oostrom. Confluence by decreasing diagrams. *Theoretical Computer Science*, 121:259–280, May 1994.
9. V. Oostrom. Developing developments. Technical Report ISRL-94-4, Basic Research Laboratories, NTT, Japan, Dec. 1994.
10. M. Oyamaguchi and Y. Toyama. On the Church-Rosser property of E-overlapping and simple-right-linear TRS's. Tech. Rep. COMP94-29 (1994-07), IEICE, 1994.
11. P. Padawitz. *Correctness, Completeness and Consistency of Equational Data Type Specifications*. PhD thesis, FB 20 (Informatik), TU Berlin, 1983. Report No. 83-15.
12. B. Rosen. Tree-manipulating systems and Church-Rosser theorems. *Journal of the ACM*, 20:160–187, 1973.
13. Y. Toyama. Commutativity of term rewriting systems. In K. Fuchi and L. Kott, eds., *Programming of Future Generation Computer*, vol. II, pp. 393–407. North-Holland, 1988.
14. Y. Toyama. Personal e-mail communication, August 31, 1995.
15. Y. Toyama and M. Oyamaguchi. Church-Rosser property and unique normal form property of non-duplicating term rewriting systems. In N. Dershowitz et al., eds., *Proc. 4th CTRS (1994)*, *LNCS* 968, pp. 316–331. Springer-Verlag, 1995.

This article was processed using the LaTeX macro package with LLNCS style